

Enhancement of the Hydrolysis Activity of F_0F_1 -ATPases Using 60 Hz Magnetic Fields

Chuanfang Chen,^{1,2} Yuanbo Cui,³ Jiachang Yue,³ Xiaolin Huo,¹ and Tao Song^{1*}

¹*Institute of Electrical Engineering, Chinese Academy of Sciences, Beijing, China*

²*Graduate School of the Chinese Academy of Sciences, Beijing, China*

³*The National Laboratory of Biomacromolecules, Institute of Biophysics, Chinese Academy of Sciences, Beijing, China*

The effects of extremely low frequency (ELF) magnetic fields on membrane F_0F_1 -ATPase activity have been studied. When the F_0F_1 -ATPase was exposed to 60 Hz magnetic fields of different magnetic intensities, 0.3 and 0.5 mT magnetic fields enhanced the hydrolysis activity, whereas 0.1 mT exposure caused no significant changes. Even if the F_0F_1 -ATPase was inhibited by *N,N*-dicyclohexylcarbodiimide, its hydrolysis activity was enhanced by a 0.5 mT 60 Hz magnetic field. Moreover, when the chromatophores which were labeled with F-DHPE were exposed to a 0.5 mT, 60 Hz magnetic field, it was found that the pH of the outer membrane of the chromatophore was unchanged, which suggested that the magnetic fields used in this work did not affect the activity of F_0 . Taken together, our results show that the effects of magnetic fields on the hydrolysis activity of the membrane F_0F_1 -ATPases were dependent on magnetic intensity and the threshold intensity is between 0.1 and 0.3 mT, and suggested that the F_1 part of F_0F_1 -ATPase may be an end-point affected by magnetic fields. *Bioelectromagnetics* 30:663–668, 2009. © 2009 Wiley-Liss, Inc.

Key words: F_0F_1 -ATPases; chromatophores; magnetic field; hydrolysis activity

INTRODUCTION

At present, the effects of extremely low frequency electromagnetic fields (ELF-EMFs) on human health have prompted considerable public concern since ELF-EMFs are ubiquitous in modern society [Repacholi and Greenebaum, 1999; Kheifets and Shimkhada, 2005; Saunders and McCaig, 2005; Paniagua et al., 2007]. The effects can be mostly attributed to the dynamics at the nucleic acid and protein levels, with possible influences on DNA, RNA, and protein synthesis [Cridland et al., 1999; Ciombor et al., 2002; Del Re et al., 2003, 2004; Schmitz et al., 2004], the membrane protein structure of living HeLa cells [Ikehara et al., 2003], ornithine decarboxylase enzyme activity [Byus, 1995; Valtersson et al., 1997; Ichinose et al., 2004; Naarala et al., 2004], Na,K-ATPase activity [Blank and Soo, 1993, 1996; Martirosov and Blank, 1995] and biological membranes [Miller, 1991; Paradisi et al., 1993; Astumian et al., 1995; Bersani et al., 1997; Volpe et al., 1998; Baureus-Koch et al., 2003]. However, there are still some disputes concerning the physical mechanism of the effects of electromagnetic fields on organisms [Adair, 1991; Valberg et al., 1997].

In a living cell the F_0F_1 -ATPases interconverts two major “energy currencies”, namely, the transmembrane electrochemical potential difference of

protons (or Na^+ in some organisms) and ATP. It is composed of two parts, membrane-embedded F_0 and hydrophilic catalytic F_1 [Kagawa et al., 1979] (Fig. 1). To further understand the effects of ELF-EMFs on human health, it is important to explore the effects of ELF-EMFs on F_0F_1 -ATPases as a possible initial step in a cascade of events that could affect health. Many reports have described the effects of electric fields on ATPase [Witt et al., 1976; Witt, 1979; Vinkler and Korenstein, 1982; Tsong, 1990; Zrimec et al., 2002], but few reports so far have focused on the effects of magnetic fields on ATPase. In fact, research into the bioeffects of ELF magnetic fields is more important than that of electric fields for evaluating health hazards

Grant sponsor: The National Natural Sciences Foundation of China; Grant number: 50377042.

*Correspondence to: Tao Song, Institute of Electrical Engineering, Chinese Academy of Sciences, Beijing 100190, China.
E-mail: songtao@mail.iee.ac.cn

Received for review 5 May 2008; Final revision received 24 November 2008

DOI 10.1002/bem.20509
Published online 3 June 2009 in Wiley InterScience (www.interscience.wiley.com).




Fig. 1. The schematic subunit organization of F_0F_1 -ATPases. F_0F_1 -ATPases is composed of eight different subunits, five of which belong to F1, subunits alpha, beta, gamma, delta, and epsilon (3:3:1:1:1), and three to F0, subunits a, b, and c (1:2:10–12).

of ELF-EMFs because there are more disputes on the bioeffects of ELF magnetic fields [Repacholi and Greenebaum, 1999]. The effects of ELF magnetic fields on membrane F_0F_1 -ATPase activity are reported in this article.

MATERIALS AND METHODS

Preparation of Chromatophores

The chromatophores were prepared from the cells of *Rhodospirillum rubrum* as previously reported [Suzuki et al., 2003; Cui et al., 2005]. In brief, *R. rubrum* was cultured anaerobically in closed vessels at 30 °C under continuous illumination according to the method described elsewhere [Zürner and Bachofen, 1979]. The harvested cells of *R. rubrum* were suspended in a buffer solution A (0.1 mM Tricine–NaOH, 0.25 M sucrose, 5 mM $MgCl_2$, pH 8.0). The suspension was incubated on ice for 30 min after addition of 1 mg/ml lysozyme. The suspension was sonicated for 10 min (5 s on, 5 s off), causing the plasma membrane to break up and invert to form vesicles with the F_0F_1 -ATPases inside out (Fig. 1). The suspension was centrifuged (25,000g, 30 min, 4 °C) and the supernatant fluid was transferred to a new tube and centrifuged again (180,000g, 90 min, 4 °C). The pellets, which are called chromatophores, were re-suspended in buffer A with 50% glycerol, frozen immediately in liquid nitrogen then stored at –80 °C.

The bacteriochlorophyll (BChl) concentration in the chromatophores was determined by the in vitro extinction coefficient as reported by Clayton [1963]. Measurements in our laboratory showed that there

were 2.34×10^{-12} mol of F_0F_1 -ATPases in 1 μ g Bchl of chromatophores.

Treatment of F_0F_1 -ATPases With DCCD and LiCl

Two procedures were used to inhibit the hydrolysis activity of F_0F_1 -ATPases. In one procedure, the F_0F_1 -ATPases were treated with *N,N*-dicyclohexylcarbodiimide (DCCD) as follows. BChl (1.2 μ g) of chromatophores was mixed with an equal volume of 100 mM Tris–HCl (pH 8.8) buffer (composed of 100 μ M DCCD) and placed on ice for 1 h. In the other procedure, the F_0F_1 -ATPases were treated with 2 M LiCl as follows. The 1.2 μ g BChl of chromatophores was centrifuged at 18,000g for 30 min and the pellet was re-suspended in buffer (250 mM sucrose, 50 mM Tricine, 4 mM $MgCl_2$, 4 mM ATP, 1 mM dithiothreitol, 1 mM phenylmethylsulfonyl fluoride, 2 M LiCl, pH 8.0) then incubated at 25 °C for 30 min to remove the beta subunits of F_0F_1 -ATPase from the chromatophores.

Hydrolysis Activity of the F_0F_1 -ATPases in Chromatophores

Liu et al. [2004] reported that when the concentration of the ATP is pre-saturated ($[ATP] = 10\text{--}100 \mu$ M) and saturated ($[ATP] > 100 \mu$ M), the hydrolysis mechanism of the F_0F_1 -ATPases behaved in an ATP-dependent manner instead of the widely believed Michaelis–Menten mechanism. In this article, the concentration of the ATP will be raised to saturation level to get good results. Accordingly, the hydrolysis activity of F_0F_1 -ATPases is the maximum rate of ATP hydrolyzed per unit of enzyme per unit time and is expressed in moles of ATP hydrolyzed per mole of F_0F_1 -ATPases per second (mol ATP/mol F_0F_1 -ATPases/s or simply 1/s).

BChl (18 μ g) of chromatophores was re-suspended in 35 μ l of ATP hydrolysis buffer (50 mM Tricine–NaOH (pH 8.0), 5 mM $MgCl_2$, 4 mM NaH_2PO_4 , 15 mM glucose, 5 mM dithiothreitol) at 25 °C. After adding 1 μ l of 10 mM ATP and mixing immediately, the F_0F_1 -ATPase hydrolysis reaction of ATP began and continued for 30 min at 25 °C. During the reaction, 2 μ l of reaction mixture was removed every 3 min and the reaction stopped using 0.2 μ l of 4% trichloroacetic acid (TCA). The ATP concentration was measured by the luciferin/luciferase system. Typically, 2.2 μ l of mixture was diluted with 100 μ l of buffer (100 mM Tricine–acetate, 2 mM EDTA, pH 7.75) and 50 μ l of luciferin/luciferase was added. The mixture was mixed quickly and placed in an ultra weak fluorescence analyzer immediately and the photons produced in 10 s at 25 °C

were recorded by computer. In this system, 1 mol ATP will produce 6.77×10^{14} photons.

Labeling the Chromatophores With pH Indicator F-DHPE

Frozen chromatophores (20 μ l) was diluted with 980 μ l buffer (0.1 mM Tricine–NaOH, 50 mM KCl, 5 mM MgCl₂, pH 8.0) and 1 μ l saturated *N*-(fluorescein-5-thiocarbonyl)-1,2-dihexadecanoyl-*sn*-glycero-3-phosphoethanolamine, triethylammonium salt added (F-DHPE) (Molecular Probes, Eugene, OR). This was then incubated on ice for 30 min and centrifuged at 15,000g for 15 min. The resulting pellet was re-suspended in buffer solution (0.1 mM Tricine–NaOH, 50 mM KCl, 5 mM MgCl₂, pH 8.0). The fluorescence of the pH indicator F-DHPE was detected by a F-4500 FL Spectrophotometer (Hitachi, Tokyo, Japan).

MF Exposure System and Experimental Design

An ELF magnetic exposure system was constructed in our laboratory for the experiments. This system consists of a high-precision AC current source and two rectangular coils, as described by Misakian et al. [1993]. Each rectangular coil was made up of two sub-coils each with 100 turns of copper wire (diameter 1 mm), which could produce equal magnetic field strength. The two rectangular coils consisted of 200 turns of copper wire with dimensions 12 cm \times 6 cm separated by a distance of 7 cm. To control the currents in the coil, a bi-directional switch was designed. When the currents in the two sub-coils were in same direction, the coil would generate and expose the sample to a uniform, sinusoidal, vertical AC magnetic field. When the currents of the sub-coils were in opposite directions, the coil would generate magnetic fields with zero strength for a sham exposure experiment. In our experiments, the exposure fields were characterized as 60 Hz, 0.1–0.5 mT, and detected by a Gauss/Tesla meter (F.W. Bell Model 7010, Bell Technologies, Orlando, FL). The non-uniformity of the magnetic fields in the exposed space (50 mm \times 10 mm \times 5 mm) is less than 2%. The vertical component of the geomagnetic field is 31 μ T, the north-south component is 28 μ T and the east–west component is 11 μ T, values which were measured by a fluxgate magnetometer (National Institute of Metrology, Beijing, P. R. China).

In this article, the exposure experiments were done in a temperature-controlled (25 ± 1 °C) room, and the temperature difference between exposure group and sham group is less than 0.2 °C in the exposed space. During the measurement, the researcher who measured the ATP concentration was not informed of the

exposure conditions. Each exposure experiment was repeated at least three times.

Statistical Analysis

All the data were analyzed using the software SPSS 13.0 for Windows (SPSS, Chicago, IL). The statistical significance was determined by a paired-sample *t*-test. Error bars represent standard deviation (SD) of the mean. The significance was set at **P* < 0.05, ***P* < 0.01.

RESULTS

Hydrolysis Activity of F₀F₁-ATPases

The hydrolysis activity of the chromatophores at room temperature (25 °C) was calculated to be 14.11 ± 1.67 s⁻¹. When the F₀F₁-ATPases were treated by the inhibitor DCCD and LiCl, the activities fell to only 4.80 ± 0.68 s⁻¹ and 0.86 ± 0.51 s⁻¹, respectively.

Effects of Magnetic Fields on the Hydrolysis Activity of F₀F₁-ATPases

The rate of autohydrolysis of ATP at 25 °C in the buffer system was $1.07 \pm 0.89 \times 10^{-13}$ mol/s. When the ATP buffer was exposed to 0.5 mT 60 Hz magnetic field, the rate was $1.17 \pm 0.95 \times 10^{-13}$ mol/s. There are no significant differences between the rates of the exposed and the sham group (*P* > 0.05, *n* = 6). Therefore, the autohydrolysis of ATP was not affected by the magnetic field.

When the membrane F₀F₁-ATPases were exposed to 60 Hz magnetic fields at different magnetic intensities (0.1, 0.3, and 0.5 mT), the results, presented in Figure 2, showed that both 0.3 and 0.5 mT, 60 Hz




Fig. 2. The effects of 0.1–0.5 mT, 60 Hz magnetic fields on F₀F₁-ATPase. Error bars represent SD of the mean. Significance was set at **P* < 0.05, ***P* < 0.01, *n* = 6.

magnetic fields enhanced the hydrolysis activity significantly (both $P < 0.05$, $n = 6$), but the 0.1 mT, 60 Hz magnetic field had no significant effect ($P > 0.05$, $n = 6$).

It is well known that the inhibitor DCCD can reduce both the synthesis and hydrolysis activity of F_0F_1 -ATPase by inhibiting the transfer of H^+ across membrane. When the F_0F_1 -ATPase had been inhibited by DCCD for 1 h, the hydrolysis activity was $4.80 \pm 0.68 s^{-1}$, which represented a 68.1% deactivation. When the F_0F_1 -ATPase, inhibited by DCCD for 1 h, was exposed to a 0.5 mT 60 Hz magnetic field, the hydrolysis activity increased to $6.97 \pm 1.24 s^{-1}$ ($P < 0.01$, $n = 6$). Therefore, exposure to a 0.5 mT 60 Hz magnetic field still stimulated the hydrolysis activity of the F_0F_1 -ATPase even after the F_0F_1 -ATPase had been inhibited by DCCD for 1 h.

When the F_0F_1 -ATPases were treated with 2 M LiCl for 24 h, the hydrolysis activity was reduced to $0.86 \pm 0.51 s^{-1}$ due to the removal of the beta subunits of F_0F_1 -ATPase. When these treated F_0F_1 -ATPases were exposed to a 0.5 mT 60 Hz magnetic field, the hydrolysis activity of F_0F_1 -ATPase was $0.66 \pm 0.34 s^{-1}$. Thus, there is no significant difference between the exposed group and the sham group ($P > 0.05$, $n = 6$).

Effects of MF on the pH of the Chromatophores

To further investigate how magnetic fields affect membrane F_0F_1 -ATPase, a H^+ transfer experiment was performed without the addition of ATP. F-DHPE is a pH indicator and has been used to measure pH changes adjacent to the bilayer surface. In the range of pH 7.0–9.0, F-DHPE is sensitive to pH changes and has a positive correlation with the change of pH [Cui et al., 2005]. When the chromatophores labeled with F-DHPE were exposed to a 0.5 mT, 60 Hz magnetic field, the fluorescence of exposed samples and the sham samples are as shown in Figure 3. Each fluorescence response curve was fitted to a linear function and the slopes calculated. It was found that the fluorescence decreased whether or not the chromatophores were exposed. However, there was no significant difference between the slopes of the fluorescence time dependence of the exposed and sham samples ($P > 0.05$, $n = 10$). This means that the pH of outer membrane of chromatophore was not changed when the chromatophores were exposed to the 0.5 mT, 60 Hz magnetic field.

DISCUSSION

The effects of ELF electromagnetic fields on membrane Na,K-ATPase have been well documented. These reports have demonstrated that the activity of




Fig. 3. The effect of 0.5 mT, 60 Hz magnetic fields on the pH of the chromatophores. The gray curve line A is fluorescence of chromatophores exposed by MF for 20 min first and then sham-exposed for the same time after 2 min interval; the dark line is linear fitting to each curve line. Curve line B was the reverse turn.

Na,K-ATPase is stimulated by magnetic fields whether the activity is high or low, and the stimulating effects decrease with increasing Na,K-ATPase activity [Blank and Soo, 1993, 1996; Martirosov and Blank, 1995]. In this article, the effects of ELF magnetic fields on membrane F_0F_1 -ATPase were stimulated by 60 Hz magnetic fields, an effect similar to the stimulating effects of magnetic fields on the activity of Na,K-ATPases. When the F_0F_1 -ATPases were exposed to 60 Hz magnetic fields of different intensities, the hydrolysis activity was enhanced by 0.3 and 0.5 mT magnetic fields but when the magnetic intensity was as low as 0.1 mT it did not show any effect. Accordingly, the effect of magnetic fields seems to depend on the magnetic intensity with the threshold lying between 0.1 and 0.3 mT.

It is well known that the DCCD is a potent inhibitor of F_0F_1 -ATPase and acts on the membrane sector, F0 [Matsuno-Yagi and Hatefi, 1993]. When the hydrolysis activity of F_0F_1 -ATPase was inactivated by 68.1% using DCCD, the 0.5 mT 60 Hz magnetic field still stimulated the hydrolysis activity of F_0F_1 -ATPase. The increase was unchanged regardless of whether the F_0F_1 -ATPase was inhibited by DCCD or not. These results suggest that the magnetic fields mainly affect F1 because DCCD acts only on F0 [Matsuno-Yagi and Hatefi, 1993].

When the chromatophores were exposed to a 0.5 mT, 60 Hz magnetic field, the pH of the outer membrane of the chromatophore was not changed. It is suggested that the magnetic fields used in this article did not affect the activity of F0 because the main function of the F0 is to transfer H^+ . Since the magnetic fields did enhance the hydrolysis activity of F_0F_1 -ATPase, it is

concluded that the effects of magnetic fields on F₀F₁-ATPase mainly result from the effect on F₁.

Many reports have described experiments showing that an external electric field can stimulate the synthesis of ATP by F₀F₁-ATPase [Witt et al., 1976; Witt, 1979; Vinkler and Korenstein, 1982; Tsong, 1990; Zrimec et al., 2002]. Vinkler and Korenstein also reported that the inhibitors Dio-9 and DCCD did not have an inhibitory effect on the ATP synthesis stimulated by external fields. They suggested that the binding site of these two inhibitors does not seem to participate in the reaction. Our results seem to be in agreement with these previous results. However, the inductive electric field in our experiment, which was computed to be at the level of $\sim 10^{-4}$ V/m, was much smaller than the external electric fields used in those articles (always greater than 100 V/m). Some reports suggested that there may be other mechanisms that respond directly to the effects of magnetic fields on F₀F₁-ATPase. Blank and Goodman [2004] suggested, from the effects of magnetic fields on Na,K-ATPase, cytochrome oxidase and the Belousov–Zhabotinski (BZ) reaction, that the responses of these simpler biological systems to magnetic fields also support a mechanism initiated by interaction with electrons. Accordingly, the description of a mechanism for the effects of magnetic fields on F₀F₁-ATPase needs further research.

It is known that there have been considerable concerns and controversies recently about the effects on human health from increasing exposure to ELF-EMFs [Merchant et al., 1994; Lacy-Hulbert et al., 1998; Saunders and McCaig, 2005; Paniagua et al., 2007]. These effects may be related to the change of activity of F₀F₁-ATPase induced by the magnetic fields, since F₀F₁-ATPase is an important protein for energy transport across membrane. For example, Sobel et al. [1996] and Qiu et al. [2004] reported that EMFs could increase the risk of Alzheimer's disease (AD) among workers. Martinez-Cano et al. [2005] reported that the hydrolytic activity of F₀F₁-ATPase increases significantly in patients with probable AD, and increased as cerebral deterioration progressed in the male population with probable AD. Since the results in this study indicate that magnetic fields affect the hydrolysis activity of F₀F₁-ATPase, it is possible to speculate that the association between magnetic fields and AD might be related to a change of the hydrolysis activity of F₀F₁-ATPase. Regardless of this speculation, the normal status of the living cell may be negatively impacted when the activity of F₀F₁-ATPase is changed by magnetic fields.

In conclusion, 60 Hz magnetic fields were found to enhance the activity of F₀F₁-ATPase. The effects depended on magnetic intensity and the magnetic fields

mainly affected the F₁. Our results may provide a new way to explore a relationship between the effects of magnetic fields on human health and the effects of magnetic fields on membrane F₀F₁-ATPase.

ACKNOWLEDGMENTS

The authors thank Dr. Yiwen Li of the Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, and Dr. Weidong Pan for his fruitful discussion. We also thank the technical staff for their work on the exposure system.

REFERENCES

- Adair RK. 1991. Constraints on biological effects of weak extremely-low-frequency electromagnetic fields. *Phys Rev A* 43(2):1039–1048.
- Astumian RD, Weaver JC, Adair RK. 1995. Rectification and signal averaging of weak electric fields by biological cells. *Proc Natl Acad Sci* 92:3740–3743.
- Baureus-Koch CL, Sommarin M, Persson BR, Salford LG, Eberhardt JL. 2003. Interaction between weak low frequency magnetic fields and cell membranes. *Bioelectromagnetics* 24(6):395–402.
- Bersani F, Marinelli F, Ognibene A, Matteucci A, Cecchi S, Santi S, Squarzone S, Maraldi NM. 1997. Intramembrane protein distribution in cell cultures is affected by 50 Hz pulsed magnetic fields. *Bioelectromagnetics* 18(7):463–469.
- Blank M, Soo L. 1993. The Na,K-ATPase as a model for electromagnetic field effects on cells. *Bioelectrochem Bioenerg* 30:85–92.
- Blank M, Soo L. 1996. Threshold for Na,K-ATPase stimulation by electromagnetic fields. *Bioelectrochem Bioenerg* 40:63–65.
- Blank M, Goodman R. 2004. Initial interactions in electromagnetic field-induced biosynthesis. *J Cell Physiol* 199:359–363.
- Byus CV. 1995. Alterations in ornithine decarboxylase activity: A cellular response to low-energy electromagnetic field exposure. In: Summary and results of radiofrequency radiation conference, Published by U.S. Environmental Protection Agency 402-R-95-011.
- Ciombor DM, Lester G, Aaron RK, Neame P, Caterson B. 2002. Low frequency EMF regulates chondrocyte differentiation and expression of matrix proteins. *J Orthop Res* 20(1):40–50.
- Clayton RK. 1963. Absorption spectra of photosynthetic bacteria and their chlorophylls. In: Gest H, Pietro AS, Vernon LP, editors. *Bacterial photosynthesis*. Antioch: Yellow Springs. pp 495–500.
- Cridland NA, Sabour NR, Saunders RD. 1999. Effects of 50 Hz magnetic field exposure on the rate of RNA synthesis by normal human fibroblasts. *Int J Radiat Biol* 75(5):647–654.
- Cui YB, Zhang F, Yue JC. 2005. Detecting proton flux across chromatophores driven by F₀F₁-ATPase using *N*-(fluorescein-5-thiocarbamoyl)-1, 2-dihexadecanoyl-*sn*-glycero-3-phosphoethanolamine, triethylammonium salt. *Anal Biochem* 344(1):102–107.
- Del Re B, Bersani F, Agostini C, Mesirca P, Giorgi G. 2004. Various effects on transposition activity and survival of *Escherichia coli* cells due to different ELF-MF signals. *Radiat Environ Biophys* 43(4):265–270.

- Del Re B, Garoia F, Mesirca P, Agostini C, Bersani F, Giorgi G. 2003. Extremely low frequency magnetic fields affect transposition activity in *Escherichia coli*. *Radiat Environ Biophys* 42(2):113–118.
- Ichinose TY, Burch JB, Noonan CW, Yost MG, Keefe TJ, Bachand A, Mandeville R, Reif JS. 2004. Immune markers and ornithine decarboxylase activity among electric utility workers. *J Occup Environ Med* 46(2):104–112.
- Ikehara T, Yamaguchi H, Hosokawa K, Miyamoto H, Aizawa K. 2003. Effects of ELF magnetic field on membrane protein structure of living HeLa cells studied by Fourier transform infrared spectroscopy. *Bioelectromagnetics* 24(7):457–464.
- Kagawa Y, Sone N, Hirata H, Yoshida M. 1979. Structure and function of H⁺-ATPase. *J Bioenerg Biomembr* 11:39–78.
- Kheifets L, Shimkhada R. 2005. Childhood leukemia and EMF: Review of the epidemiologic evidence. *Bioelectromagnetics* 26(Suppl. 7):S51–S59.
- Lacy-Hulbert A, Metcalfe JC, Hesketh R. 1998. Biological responses to electromagnetic fields. *Fed Am Soc Exp Biol J* 12(6):395–420.
- Liu MS, Todd BD, Sados RJ. 2004. Complex cooperativity of ATP hydrolysis in the F₁-ATPase molecular motor. *Biochim Biophys Acta* 1752:111–123.
- Martinez-Cano E, Ortiz-Genaro G, Pacheco-Moises F, Macias-Islas MA, Sanchez-Nieto S, Rosales-Corral SA. 2005. Functional disorders of F₀F₁-ATPase in submitochondrial particles obtained from platelets of patients with a diagnosis of probable Alzheimer's disease. *Rev Neurol* 40(2):81–85.
- Martirosov S, Blank M. 1995. Inhibition of F₀F₁-ATPase activity in AC-fields. *Bioelectrochem Bioenerg* 37:153–156.
- Matsuno-Yagi A, Hatefi Y. 1993. Studies on the mechanism of oxidative phosphorylation. ATP synthesis by submitochondrial particles inhibited at F₀ by venturicidin and organotin compounds. *J Biol Chem* 268(9):6168–6173.
- Merchant CJ, Renew DC, Swanson J. 1994. Exposure to power-frequency magnetic fields in the home. *J Radiol Protect* 14:77–87.
- Miller DL. 1991. Electric fields induced in chicken eggs by 60-Hz magnetic fields and the dosimetric importance of biological membranes. *Bioelectromagnetics* 12(6):349–360.
- Misakian M, Sheppard AR, Krause D, Frazier ME, Miller DL. 1993. Biological, physical, and electrical parameters for in vitro studies with ELF magnetic and electric fields: A primer. *Bioelectromagnetics* 14(Suppl 2):1–73.
- Naarala J, Hoyto A, Markkanen A. 2004. Cellular effects of electromagnetic fields. *Altern Lab Anim* 32(4):355–360.
- Paniagua JM, Jiménez A, Rufo M, Gutiérrez JA, Gómez FJ, Antolín A. 2007. Exposure to extremely low frequency magnetic fields in an urban area. *Radiat Environ Biophys* 46(1):69–76.
- Paradisi S, Donelli G, Santini MT, Straface E, Malorni W. 1993. A 50 Hz magnetic field induces structural and biophysical changes in membranes. *Bioelectromagnetics* 14(3):247–255.
- Qiu C, Fratiglioni L, Karp A, Winblad B, Bellander T. 2004. Occupational exposure to electromagnetic fields and risk of Alzheimer's disease. *Epidemiology* 15(6):687–694.
- Repacholi MH, Greenebaum B. 1999. Interaction of static and extremely low frequency electric and magnetic fields with living systems: Health effects and research needs. *Bioelectromagnetics* 20(3):133–160.
- Saunders RD, McCaig CD. 2005. Developmental effects of physiologically weak electric fields and heat: An overview. *Bioelectromagnetics* 26(7 Suppl):S127–S132.
- Schmitz C, Keller E, Freuding T, Silny J, Korr H. 2004. 50-Hz magnetic field exposure influences DNA repair and mitochondrial DNA synthesis of distinct cell types in brain and kidney of adult mice. *Acta Neuropathol (Berl)* 107(3):257–264.
- Sobel E, Dunn M, Davanipour Z, Qian Z, Chui HC. 1996. Elevated risk of Alzheimer's disease among workers with likely electromagnetic field exposure. *Neurology* 47(6):1477–1481.
- Suzuki T, Murakami T, Iion R, Suzuki J, Ono S, Shirakihara Y, Yoshida M. 2003. F₀F₁-ATPase/synthase is geared to the synthesis mode by conformational rearrangement of ε subunit in response to proton motive force and ADP/ATP ratio. *J Biol Chem* 278:46840–46846.
- Tsong TY. 1990. Electrical modulation of membrane proteins: Enforced conformational oscillations and biological energy and signal transductions. *Annu Rev Biophys Chem* 19:83–106.
- Valberg PA, Kavet R, Rafferty CN. 1997. Can low-level 50/60 Hz electric and magnetic fields cause biological effects? *Radiat Res* 148(1):2–21.
- Valtersson U, Mild KH, Mattsson MO. 1997. Ornithine decarboxylase activity and polyamine levels are different in Jurkat and CEM-CM3 cells after exposure to a 50 Hz magnetic field. *Bioelectrochem Bioenerg* 43:169–172.
- Vinkler C, Korenstein R. 1982. Characterization of external electric field-driven ATP synthesis in chloroplasts. *Proc Natl Acad Sci USA* 79(10):3183–3187.
- Volpe P, Parasassi T, Esposito C, Ravagnan G, Giusti AM, Pasquarelli A, Eremenko T. 1998. Cell membrane lipid molecular dynamics in a solenoid versus a magnetically shielded room. *Bioelectromagnetics* 19:107–111.
- Witt HT, Schlodder E, Gräber P. 1976. Membrane-bound ATP synthesis generated by an external electrical field. *FEBS Lett* 69(1):272–276.
- Witt HT. 1979. Energy conversion in the functional membrane of photosynthesis. Analysis by light pulse and electric pulse methods. *Biochim Biophys Acta* 505(3–4):355–427.
- Zrimec A, Jerman I, Lahajnar G. 2002. Alternating electric fields stimulate ATP synthesis in *Escherichia coli*. *Cell Mol Biol Lett* 7:172–174.
- Zürner H, Bachofen R. 1979. Hydrogen production by the photosynthetic bacterium *rhodospirillum rubrum*. *Appl Environ Microbiol* 37(5):789–793.