

# Direct differentiation of atrial and ventricular myocytes from human embryonic stem cells by alternating retinoid signals

Qiangzhe Zhang<sup>1,\*</sup>, Junjie Jiang<sup>1,\*</sup>, Pengcheng Han<sup>1,\*</sup>, Qi Yuan<sup>1</sup>, Jing Zhang<sup>1</sup>, Xiaoqian Zhang<sup>1</sup>, Yanyan Xu<sup>1</sup>, Henghua Cao<sup>1</sup>, Qingzhang Meng<sup>1</sup>, Li Chen<sup>1</sup>, Tian Tian<sup>1</sup>, Xin Wang<sup>2</sup>, Pu Li<sup>1</sup>, Jurgen Hescheler<sup>3</sup>, Guangju Ji<sup>1</sup>, Yue Ma<sup>1</sup>

<sup>1</sup>National Laboratory of Biomacromolecules, Institute of Biophysics, Chinese Academy of Sciences, 15 Datun Rd., Chaoyang District, Beijing 100101, China; <sup>2</sup>Institute of Biochemistry and Cell Biology, Shanghai Institutes of Biological Sciences, 320 Yueyang Rd., Shanghai 200032, China; <sup>3</sup>Center for Physiology and Pathophysiology, Institute for Neurophysiology, University of Cologne, Robert-Koch-Str. 39, 50931 Cologne, Germany

Although myocyte cell transplantation studies have suggested a promising therapeutic potential for myocardial infarction, a major obstacle to the development of clinical therapies for myocardial repair is the difficulties associated with obtaining relatively homogeneous ventricular myocytes for transplantation. Human embryonic stem cells (hESCs) are a promising source of cardiomyocytes. Here we report that retinoid signaling regulates the fate specification of atrial versus ventricular myocytes during cardiac differentiation of hESCs. We found that both Noggin and the pan-retinoic acid receptor antagonist BMS-189453 (RAi) significantly increased the cardiac differentiation efficiency of hESCs. To investigate retinoid functions, we compared Noggin+RAi-treated cultures with Noggin+RA-treated cultures. Our results showed that the expression levels of the ventricular-specific gene *IRX-4* were radically elevated in Noggin+RAi-treated cultures. *MLC-2V*, another ventricular-specific marker, was expressed in the majority of the cardiomyocytes in Noggin+RAi-treated cultures, but not in the cardiomyocytes of Noggin+RA-treated cultures. Flow cytometry analysis and electrophysiological studies indicated that with  $64.7 \pm 0.88\%$  (mean  $\pm$  s.e.m) cardiac differentiation efficiency, 83% of the cardiomyocytes in Noggin+RAi-treated cultures had embryonic ventricular-like action potentials (APs). With  $50.7 \pm 1.76\%$  cardiac differentiation efficiency, 94% of the cardiomyocytes in Noggin+RA-treated cultures had embryonic atrial-like APs. These results were further confirmed by imaging studies that assessed the patterns and properties of the  $Ca^{2+}$  sparks of the cardiomyocytes from the two cultures. These findings demonstrate that retinoid signaling specifies the atrial versus ventricular differentiation of hESCs. This study also shows that relatively homogeneous embryonic atrial- and ventricular-like myocyte populations can be efficiently derived from hESCs by specifically regulating Noggin and retinoid signals.

**Keywords:** human embryonic stem cell; cardiac subtype specification; heart development

*Cell Research* (2011) 21:579-587. doi:10.1038/cr.2010.163; published online 23 November 2010

## Introduction

Using current technologies, embryonic atrial-, ventric-

ular- and nodal-like cardiomyocytes can be differentiated nonspecifically from hESCs [1-4]. Swine transplantation studies have shown that implanted hESC-derived cardiomyocytes have pace-making activities, which are a potential cause of ventricular arrhythmias [5]. The application of hESCs in myocardial repair is hampered by this cardiac subtype heterogeneity of hESC-derived cardiomyocytes [6]. To direct the differentiation of hESCs into a desired cardiac subtype, the mechanisms of cardiac subtype specification must be uncovered. Although sev-

\*These three authors contributed equally to this work.

Correspondence: Yue Ma<sup>a</sup>, Guangju Ji<sup>b</sup>

<sup>a</sup>E-mail: yuema@ibp.ac.cn

<sup>b</sup>E-mail: gj28@ibp.ac.cn

Received 30 September 2010; revised 3 November 2010; accepted 8 November 2010; published online 23 November 2010

eral growth factors, such as activin A, bone morphogenetic protein 4 (BMP4), wnt-3a, basic fibroblast growth factor (bFGF) and dickkopf homolog 1 (DKK1), promote cardiogenesis and are used in several hESC cardiac differentiation protocols [2, 3, 7], there is no evidence to date showing that these or other growth factors regulate cardiac subtype specification during hESC differentiation. Identifying the key regulators of cardiac subtype specification is critical for reducing the heterogeneity of the hESC-derived cardiomyocyte population. Reducing the heterogeneity of this population will be important for its later use in regenerative medicine or in drug testing systems [1, 8].

BMP signaling is tightly controlled during mesoderm and heart development. In mouse embryos, the BMP antagonist Noggin is transiently but strongly expressed in the cardiac crescent at embryonic day E7.5 to E8.0 [9]. Dkk1, a potent inducer of heart development [3, 10, 11], synergises with BMP antagonism to specify heart tissue in non-cardiogenic mesoderm from *Xenopus* embryos [12]. It has been shown that long-term treatment of hESCs with BMP4 induces trophoblast-like cell differentiation [13], while short-term treatment initiates mesoderm formation [14]. Together, these results suggest that inhibiting BMP signaling after mesoderm formation facilitates cardiac development.

RA signaling restricts the cardiac progenitor pool, and exposure of the anterior lateral plate mesoderm of zebrafish embryos to the RA antagonist BMS-189453 causes uncommitted lateral mesodermal cells to become myocardial progenitors [15, 16]. RA signaling also regulates anterior-posterior polarisation of the heart [17]. Chicken transplantation studies have revealed that the cardiogenic mesoderm from HH stages 4-6, originally fated to be atria, is competent to develop into functional ventricles, and vice versa [18, 19]. RA treatment of HH stage 4 cardiogenic tissue activates the expression of the atrium-specific gene *AMHCI* in anterior progenitors fated to develop into out-flow track tissues [20]. Furthermore, in both mouse and chicken embryos, inhibition of RA signaling within critical periods produces embryos with oversized ventricles and smaller or missing atria, and the exogenous addition of RA results in reverted phenotypes [21, 22]. Furthermore, studies with mouse embryonic stem cells indicate that retinoic acid promotes the expression of atrial-related genes [23].

Based on these previous studies, we hypothesised that inhibition of the BMP pathway after the initiation of hESC differentiation and blocking retinoic acid signaling would promote cardiogenesis. Retinoid signaling may also regulate the atrial versus ventricular differentiation of hESCs. To test these hypotheses, we added Noggin,

RA and the RA inhibitor RAi to cardiac differentiation cultures at different time intervals and investigated the effects of these factors on the cardiogenesis and cardiac subtype specification of hESC derivatives. Our results indicate that the inhibition of both BMP and RA signals with Noggin and RAi significantly promotes cardiogenesis, and retinoid signaling controls the atrial versus ventricular specification of differentiated hESCs. In addition to providing important insights into the mechanisms that specify cardiac subtypes, our findings also demonstrate the direct differentiation of relatively homogeneous embryonic atrial- and ventricular-like myocytes from hESCs.

## Results

### *Noggin and the RA antagonist BMS-189453 promote the cardiogenesis of differentiated hESCs*

To investigate its functions in cardiac differentiation, Noggin was systematically added at different time intervals between days 2 and 5 to cardiomyocyte-differentiating hESC cultures generated by a protocol developed in our laboratory (see Methods for a detailed description). The results show that cardiac differentiation was slightly repressed when Noggin was present between days 2 and 3, while differentiation was significantly promoted when Noggin was present between days 2.5 and 4.5. The highest cardiac differentiation efficiency was achieved when Noggin was applied between days 4 and 5 (Figure 1B). Western blotting for phosphorylated Smad 1, 5 and 8 indicated that Noggin reduced the downstream activities of BMP signaling (data not shown). Therefore, the inhibition of BMP signaling promotes cardiogenesis in hESCs after the initiation of differentiation.

Previous reports that RA signaling restricts the embryonic cardiac progenitor pool raise the possibility that inhibition of RA signaling during cardiac differentiation of hESCs could promote cardiogenesis. Vitamin A, the substrate for RA synthesis, and RALDH2, the enzyme responsible for RA synthesis [21], were both present in our cultures (data not shown), suggesting that RA signaling was potentially activated. Therefore, we tested the effects of RA inhibition on hESC cardiac differentiation by adding RAi to our cardiomyocyte-differentiating cultures between days 4 and 9 (Figure 1C). Flow cytometry analysis indicated that cardiac differentiation efficiencies were markedly increased when RAi was added between days 6 and 9 (Figure 1C). These results demonstrated that inhibition of RA signaling promotes cardiac differentiation of hESCs.


Next, we combined Noggin treatment on days 4 and 5 with RAi treatment on days 6, 7 and 8. Flow cytometry

analysis of CTNT<sup>+</sup> cells from day 14 cultures showed that the differentiation efficiency with Noggin alone was 50 ± 3.06% (mean ± s.e.m.), and this efficiency increased to 73 ± 2.08% when the cells were treated with both RAi and Noggin (Figure 1D). This result was confirmed by quantitative RT-PCR analysis of day 14 cultures. The expression levels of both CTNT and NKX2.5 were significantly higher in the Noggin+RAi-treated cultures than in cultures treated with Noggin alone (Figure 1E). Immunostaining indicated the expression of typical cardiac markers, including CTNT,  $\alpha$ -Actinin, MLC-2A, MLC-2V, and  $\beta$ -MHC, in these cultured cells (Figure 1F).

*Alternative retinoid signals direct the differentiation of hESCs into two distinct subtypes of cardiomyocytes*

Chicken and mouse embryo studies indicated that retinoid signaling regulates the fate specification of in-flow and out-flow track tissues [17-22]. Therefore, we hypothesized that the activation or inactivation of retinoid signaling directs the atrial vs. ventricular fate specification of differentiated hESC cardiac progenitors, and such a mechanism could be used to efficiently generate either hESC-derived atrial- or ventricular-like myocytes.


To test this hypothesis, either RA or its antagonist, RAi, was added to the Noggin-treated cultures between days 6 and 8 in parallel experiments (Figure 1A). After 14 days of differentiation, the frequencies of CTNT<sup>+</sup> cells in the Noggin+RA- and Noggin+RAi-treated cultures were 50.7 ± 1.76% and 64.7 ± 0.88%, respectively (Figure 2A). While there was only a 14% difference in the




**Figure 1** Noggin (Ngn) and RA inhibition promote hESC cardiac differentiation. **(A)** An outline of the protocol used for the differentiation of hESCs to cardiac lineages. **(B)** Frequencies of CTNT<sup>+</sup> cells at day 14; cardiac-induced cultures with addition of Ngn at the time intervals indicated; the basic protocol (BP) was used as a control. **(C)** Frequencies of CTNT<sup>+</sup> cells in day 14 cardiac-induced cultures with addition of the RA inhibitor, RAi, at the indicated time intervals; the BP was used as a control. **(D)** Flow cytometry analysis of CTNT<sup>+</sup> cells in cultures with the BP used as a control; BP cultures with Ngn added at days 4 and 5, and BP cultures with Ngn added at days 4 and 5 and RAi added from day 6 to day 8 (Ngn+RAi). **(E)** Quantitative RT-PCR gene expression analysis on day 14 BP, Ngn-treated, and Ngn+RAi-treated cultures, as indicated in **(D)**. The average expression, normalized to GADPH, is shown. **(F)** Immunostaining analysis of cardiac-induced cultures. Where shown, bars represent the standard error of the mean of three independent experiments. CM, conditioned medium. *P* compared with the BP control or otherwise indicated: \**P* < 0.05, \*\**P* < 0.005, \*\*\**P* < 0.0005.

differentiation efficiencies, the size of the beating cardiomyocytes in the Noggin+RA-treated cultures was smaller than that of those in the Noggin+RAi-treated cultures (Figure 2B and 2D). The beating rate of the cardiomyocytes in the Noggin+RA-treated cultures was also faster than that of those in the Noggin+RAi-treated cultures (Figure 2C and Supplementary information, Movies S1, S2), suggesting that there were two different subtypes of cardiomyocytes present in these two different cultures.

Next, we examined the expression of ventricular-



**Figure 2** Morphologies and beating rates of differentiated cardiomyocytes. **(A)** Flow cytometry analysis of CTNT<sup>+</sup> cells from day 14 cultures differentiated with the BP plus treatments with Ngn at days 4 and 5, and RA (Ngn+RA) or RAi (Ngn+RAi) treatment from days 6 to 8. **(B)** The size of single cardiomyocytes from 60-day-old cultures treated with Ngn+RA ( $n = 35$ ) and Ngn+RAi ( $n = 31$ ), measured with ImageJ software. **(C)** The beating rates (beat/minute) of 60-day-old differentiated cultures treated with Ngn+RA or Ngn+RAi ( $n = 4$ ). **(D)** Immunostaining of single cells from 60-day-old differentiated cultures treated with Ngn+RA or Ngn+RAi. **\*\*** $P < 0.005$ , **\*\*\*** $P < 0.0005$ .



**Figure 3** Characterization of hESC-derived cardiomyocytes induced by Ngn+RA and Ngn+RAi. **(A)** Quantitative RT-PCR analysis of the kinetics of *IRX4* gene expression in Ngn+RA and Ngn+RAi treated cultures. The average expression, normalized to GAPDH, is shown. **(B)** Immunostaining of 60-day-old Ngn+RA- and Ngn+RAi-induced cultures demonstrated MLC-2V expression in the majority of CTNT<sup>+</sup> cells of Ngn+RAi-treated cultures but not in those of Ngn+RA-treated cultures. **(C)** Western blotting of 60-day-old Ngn+RA- and Ngn+RAi-treated cultures indicated that even though CTNT is evenly expressed in both cultures, MLC-2V is strongly expressed in Ngn+RAi-treated cultures, but not in Ngn+RA-treated cultures. **\*\*\*** $P < 0.0005$ .

specific genes (*IRX4* and *MLC-2V*) in the two cultures [22, 24, 25]. Quantitative RT-PCR showed that in the Noggin+RAi-treated cultures, *IRX4* expression started to rise at day 8, and it was 10-fold greater by day 14 than the Noggin+RA-treated cultures (Figure 3A). Immunostaining of 60-day-old cultures showed that MLC-2V was expressed in the majority of CTNT<sup>+</sup> cells in the Noggin+RAi-treated cultures, but not in the Noggin+RA-treated cultures (Figure 3B). This result is consistent with the results from western blotting experiments that


indicated that although CTNT was expressed at similar levels in these two cultures, MLC-2V was strongly and only expressed in the Noggin+RAi-treated cultures (Figure 3C).

We also compared the expression levels of CTNT and MLC-2V in Noggin+RAi-, Noggin- and Noggin+RA-treated cultures by immunostaining and western blots. The results show that in the Noggin-only-treated cultures, only approximately 35% of the CTNT<sup>+</sup> cells were also MLC-2V positive, with weak MLC-2V expression detected by western blots (Supplementary information, Figure S1). Collectively our results indicate that the majority of the cardiomyocytes in the Noggin+RAi-treated cultures are embryonic ventricular-like myocytes, while the cardiomyocytes differentiated in the Noggin+RA-treated cultures are either embryonic nodal- or atrial-like myocytes that do not express MLC-2V. We also examined the expression levels of  $\beta$ -MHC, MLC-2A, and atrial natriuretic factor (ANF) in RA- and RAi-treated cultures by western blotting, and the results showed that  $\beta$ -MHC is evenly expressed in the two cultures. How-

ever, MLC-2A and ANF are expressed at higher levels in the Noggin+RA-treated cultures compared to the Noggin+RAi-treated cultures (Supplementary information, Figure S2).

*Electrophysiological characterization identifies embryonic atrial- and ventricular-like cardiomyocyte populations induced by alternative retinoid signals*

Due to a lack of endogenous early atrium-specific genetic markers in mammalian systems [22], we chose to use electrophysiological characteristics to rigorously identify these two cardiac sub-populations. Based on the morphology and classification of AP properties (Table 1) [1, 26], three major types of AP (nodal-like, atrial-like, and ventricular-like) were observed in our study (Figure 4A). However, the ratios of the three major types of AP were different between the Noggin+RA- and Noggin+RAi-treated cultures. In the cultures treated with Noggin+RAi, 83% of myocytes ( $n = 23$ ) possessed ventricular-like APs (Figure 4A and 4C), and the duration of the APs could be shortened by the application of the cal-



**Figure 4** AP morphologies and Ca<sup>2+</sup> release properties of cardiomyocytes induced by Ngn+RA and Ngn+RAi. **(A)** Three major types of APs were observed in hESC-derived cardiomyocytes: nodal-like, atrial-like, and ventricular-like. The duration of the ventricular-like APs was greatly reduced by the application of nifedipine **(B, left)**; no significant effect of nifedipine on atrial-like AP duration was observed **(B, right)**. **(C)** The percentages of the AP types recorded from Ngn+RA- and Ngn+RAi-treated cultures are indicated. **(D)** Typical Ca<sup>2+</sup> images recorded from cardiomyocytes in Ngn+RA cultures (left) and Ngn+RAi cultures (right). Fluorescence profiles (bottom) were taken from the images (at positions indicated by arrows). The properties of typical Ca<sup>2+</sup>-release events recorded from cardiomyocytes in Ngn+RA- and Ngn+RAi-treated cultures are summarized in **(E)** (98 sparks from 18 cells in Ngn+RA-treated cultures and 348 sparks from 14 cells in Ngn+RAi-treated cultures were measured). \* $P < 0.05$ . All cells used in electrophysiological studies were 60 to 90 days old. N-like, nodal-like AP; A-like, atrial-like AP; V-like, ventricular-like AP.

**Table 1** AP parameters recorded from hESC-derived cardiomyocytes

	n(cell)	Vmax(V/s)	APA(mV)	APD90(ms)	MDP(mV)
Nodal-like	6	7.3 ± 4.2	74.5 ± 8.9 <sup>#</sup>	147.6 ± 26.8 <sup>S</sup>	-51.3 ± 8.2 <sup>†</sup>
Atrial-like	36	12.8 ± 3.1*	81.6 ± 11.5 <sup>#</sup>	168.8 ± 26.8 <sup>S</sup>	-55.5 ± 6.5 <sup>†</sup>
Ventricular-like	42	11.4 ± 2.8*	86.8 ± 12.4 <sup>#</sup>	285.8 ± 52.6 <sup>S</sup>	-62.3 ± 8.6 <sup>†</sup>

Data are means ± se. n indicates the number of cells tested. Vmax, maximum rate of AP increase; APA, AP amplitude; APD90, AP duration measured at 90% repolarisation; MDP, maximum diastolic potential. \**P* < 0.05 compared with nodal-like; <sup>#</sup>*P* < 0.05 compared with each other; <sup>S</sup>*P* < 0.01 compared with each other; and <sup>†</sup>*P* < 0.05 compared with each other.

cium channel blocker nifedipine (Figure 4B, left). However, 94% of the myocytes (*n* = 19) from Noggin+RA-treated cultures exhibited an atrial-like AP, and the duration of the AP could not be shortened by nifedipine (Figures 4A, 4B right and 4C). These results demonstrate that the majority of the cardiomyocytes in the Noggin+RA-treated cultures were embryonic atrial-like myocytes, and the majority of those in the Noggin+RAi-treated cultures were embryonic ventricular-like myocytes. Interestingly, in both the Noggin+RA and Noggin+RAi-treated cultures, we did not observe the high percentages of cardiomyocytes bearing nodal-like APs that were reported in previous studies [1, 4].

There are important kinetic differences in Ca<sup>2+</sup> sparks, the elementary unit of cardiomyocyte Ca<sup>2+</sup> signaling, in atrial versus ventricular myocytes [27, 28]. Ca<sup>2+</sup> sparks are significantly larger and longer lasting in atrial myocytes compared to ventricular myocytes [28, 29]. The results of imaging studies indicated that in the Noggin+RAi-treated culture, 87.5% (14/16) of the cells tested displayed Ca<sup>2+</sup> sparks with a relatively low amplitude, fast rise time, short half time decay and small size. These are typical Ca<sup>2+</sup> spark properties of ventricular-like myocytes (Figures 4D and 4E). On the other hand, in Noggin+RA-treated cultures, 81.8% (18/22) of the cells tested displayed Ca<sup>2+</sup> sparks with a higher amplitude, slower rise time, longer half time decay and larger size (Figure 4D and 4E), suggesting that the myocytes from Noggin+RA-treated cultures were atrial-like myocytes. The kinetic study of Ca<sup>2+</sup> release and the ratios of cardiomyocytes bearing those two patterns in the two cultures are consistent with and support previous AP phenotyping-based cardiac subtype categorization.

## Discussion

Our results show that inhibition of BMP signaling after the initiation of cardiac differentiation promotes the cardiogenesis of hESCs. This finding is partially consistent with studies on mouse embryonic stem cells (mESCs) that demonstrate that the administration of Noggin before

the initiation of differentiation promotes cardiogenesis [9]. Further comparing Noggin-treated cells with non-treated cells revealed that granulocyte colony-stimulating factor (G-CSF) promotes the proliferation of developing cardiomyocytes derived from mESCs [30]. Noggin sustains the undifferentiated proliferation of hESCs [31], and BMP4 is required for mESC self-renewal [32]. These different self-renewal mechanisms may be the cause of the differences observed in the cardiac differentiation studies of human and mouse embryonic stem cells.

Even though western blots revealed the differential expression of ANF and MLC-2A, both genes are expressed in 60-day-old Noggin+RA- and Noggin+RAi-treated cultures (Supplementary information, Figure S2). This result is consistent with Dr Rosenthal's statement, which indicated that there is no early atrial-specific marker in the mouse system. Instead, they used a proximal 840-bp quail SMYHC3 promoter to label the sino-atrial tissue from the earliest stage of heart development [22].

Previous studies of chicken and mouse embryos have proposed that RA signaling determines sinoatrial cell fate, whereas ventricular fate is specified in the absence of RA [17]. Our study shows that blocking RA signaling induces the expression of the ventricle-specific marker MLC-2V in major hESC-derived cardiomyocytes, and these cells possess APs and Ca<sup>2+</sup> sparks typical of ventricular myocytes. Exogenous RA treatment directs the differentiation of hESCs into myocytes that retain the characteristic atrial-like APs and large Ca<sup>2+</sup> sparks or Ca<sup>2+</sup> transients. Our results demonstrate that the activation or inhibition of retinoid signals instructs the atrial versus ventricular specification of differentiating hESCs, respectively. By contrast, a previous study demonstrated that RA enhances the development of ventricular cardiomyocytes derived from mESCs [33]. This result could represent differences between the differentiation culture systems used in the two studies, the embryoid body procedure versus the flat culture system, or the timing of RA administration.

The potential risk of ventricular arrhythmias caused by the heterogeneity of hESC-derived cardiomyocytes

is one of the major hurdles for the application of hESCs to cardiac repair [1, 5, 6]. The application of relatively homogeneous ventricular myocytes derived from hESCs in myocardial repair has a great potential to reduce this risk by removing one of the major barriers for developing hESC-based myocardial repair strategies. Recent advances in tissue engineering, including mouse ventricular progenitor cell isolation from mESC derivatives using a genetic labelling approach to generate functional ventricular muscles [34], suggest that human functional ventricular heart muscles can be generated with embryonic ventricular-like myocytes that are directly differentiated from hESCs. With a chemically defined culture system and no genetic manipulation, the direct differentiation procedures that we have developed could be easily employed in clinical studies of myocardial repair. Another challenge for developing hESC-based myocardial repair strategies is the development of biotechnology to rapidly generate the large amounts of ventricular myocytes required for transplantation. Our study demonstrated the efficient differentiation of embryonic atrial- and ventricular-like myocytes from hESCs by eliminating the embryoid body procedure, a time-consuming step commonly used in the cardiac differentiation of embryonic stem cells. If combined together with induced pluripotent stem (iPS) cell technology [35, 36], the programmed differentiation of atrial- and ventricular-like myocytes may not only be used to develop safe cell sources for personalised cardiac repair but also provide cellular models for the study of genetic atrial or ventricular diseases.

## Materials and Methods

### *Maintenance and differentiation of hESCs*

The undifferentiated hESC line H7 (WiCell Research Institute) was maintained on matrigel-coated plates, as previously described [37]. In the basic cardiac induction protocol (BP), undifferentiated hESCs were seeded on gelatin-coated plates at a density of  $1.5 \times 10^5$  cells/cm<sup>2</sup> and cultured with mouse embryonic fibroblast-conditioned medium for 3 days until fully confluent. To initiate cell differentiation, the medium was changed to RPMI1640 and supplemented with B27 (Invitrogen). The cells were treated with 25 ng/ml BMP4 and 6 ng/ml bFGF at day 1 100 ng/ml activin A at day 2, and 200 ng/ml DKK1 (R&D Systems) from day 6 to day 11. The medium was changed every 3 days after day 11 (Figure 1). Noggin (250 ng/ml), RA (1  $\mu$ M, Sigma) or RAi (1  $\mu$ M) were added to the cell cultures at the times specified in Figures 1A-1C. Spontaneous beating clusters were typically observed on days 10 and 11. Cardiac differentiation efficiency was analysed on day 14 with CTNT antibody staining and flow cytometry analysis.

### *Single-cell preparation of hESC-derived cardiomyocytes*

Differentiated cultures that were 60 to 90 days old were washed in a low Ca<sup>2+</sup> solution and then incubated in an enzyme solution for 20 min at 37 °C. The dissociation was completed in KB

solution by gently shaking for 40 min at room temperature. The isolated cells were resuspended in DMEM plus 10% FBS, transferred to 0.1% gelatin-coated glass coverslips, and then kept in an incubator at 37 °C and 5% CO<sub>2</sub>. The composition of the low-Ca<sup>2+</sup> solution was as follows: 120 mM NaCl, 5.4 mM KCl, 5 mM MgSO<sub>4</sub>, 5 mM Na pyruvate, 20 mM glucose, 20 mM taurine, and 10 mM HEPES. The pH was adjusted to 7.3 with NaOH. The KB solution contained 85 mM KCl, 30 mM K<sub>2</sub>HPO<sub>4</sub>, 5 mM MgSO<sub>4</sub>, 1 mM EGTA, 2 mM Na<sub>2</sub>ATP, 5 mM Na pyruvate, 20 mM glucose, 20 mM taurine, and 5 mM creatine. The pH was adjusted to 7.3 with KOH.

### *Electrophysiological measurements and confocal Ca<sup>2+</sup> imaging*

The APs of cardiomyocytes were recorded in a whole-cell patch clamp configuration using an Axon 200B amplifier (Axon Instruments) at room temperature. Data were digitised at 20 kHz, filtered at 2 kHz, and analysed by the PClamp 9.0 program. Patch pipettes (2-4 M $\Omega$  resistance) were filled with an intracellular solution containing 50 mM KCl, 60 mM K-aspartate, 1 mM MgCl<sub>2</sub>, 3 mM Na<sub>2</sub>ATP, 10 mM EGTA, and 10 mM HEPES. The pH was adjusted to 7.3 with KOH. Normal Tyrode's solution was used as an extracellular solution and contained 140 mM NaCl, 5 mM KCl, 1 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub>, 10 mM glucose, and 10 mM HEPES. The pH was adjusted to 7.4 with NaOH.

For Ca<sup>2+</sup> confocal imaging, myocytes were incubated with Fluo-4AM (10  $\mu$ M/L; Molecular Probes) for 10 min at room temperature and then perfused with extracellular buffer for approximately 30 min. Ca<sup>2+</sup> imaging studies were performed on a Leica SP5 confocal microscope equipped with an argon laser (488 nm) at a magnification of 40 $\times$  using a 1.25 NA oil immersion objective. Spontaneous Ca<sup>2+</sup> sparks and Ca<sup>2+</sup> transients were recorded using linescans that were obtained at 0.5 ms per line. Images were processed and analysed using both MATLAB 7.1 software (MathWorks) and ImageJ (Scioncorp). Detection criteria of 3.8 $\times$  SD for Ca<sup>2+</sup> sparks were set, and automated counting of Ca<sup>2+</sup> sparks was performed using the Sparkmaster plug-in for ImageJ [38].

### *Flow cytometry*

Differentiated cell clusters were dissociated into single cells with 0.25% trypsin-EDTA and were then fixed and stained with anti-human CTNT antibody (R&D Systems) and goat anti-mouse FITC-conjugated secondary antibody (Santa Cruz) in PBS plus 0.5% BSA and 0.1% saponin (Sigma) at 4 °C. Stained cells were kept in 4% paraformaldehyde for subsequent quantitative analysis. Data were collected using a FACScalibur (Becton Dickinson) and analysed with FlowJo software (Treestar).

### *Real-time RT-PCR*

Total RNA was isolated from a single well of a 24-well plate of differentiated hESCs using Qiagen's RNeasy Plus Mini kit. Then, 1  $\mu$ g of total RNA was reverse transcribed with the SuperScript III First-Strand Synthesis System (Invitrogen). RT-PCR was performed using rTaq DNA Polymerase (Takara). Real-time PCR was performed in triplicate using the 2 $\times$  QuantiFast SYBR Green I PCR Master Mix (Qiagen) on a Rotor Gene 6200 Real-Time PCR Machine (Corbett) with an annealing temperature of 60 °C. The expression of each gene was normalized to GAPDH gene expression. Primer sequences are listed in Table 2.

**Table 2** Primer sequences used for qPCR

Gene	Forward Primer	Reverse Primer	Tm
<i>NXK2.5</i>	acctcaacagctccctgactct	ataatgccgcccacaaactctcc	60 °C
<i>CTNT</i>	ttaccaaaagatctgctcctcgct	ttattactggtgtggagtgggtgtgg	60 °C
<i>IRX4</i>	ttccgttctgaagcgtggtc	tgaagcaggcaattattggtgt	60 °C
<i>GAPDH</i>	gaaatcccatcaccatctccagg	gagcccccagccttctccatg	60 °C

### Immunofluorescence

Differentiated cultures (60 days old) were digested with 0.25% trypsin-EDTA, and the cells were plated on gelatin-coated coverslips for 5 days to allow full attachment to occur. Cells were then fixed in 4% paraformaldehyde and incubated with primary antibodies: anti-human CTNT (R&D systems), mouse anti-human  $\alpha$ -Actinin (Sigma), mouse anti-human  $\beta$ -MHC (ATCC), mouse anti-human MLC-2A (Synaptic Systems), or rabbit anti-human MLC-2V (ProteinTech Group). Goat anti-mouse secondary antibody conjugated with DyLight 488 (Santa Cruz Biotechnology) and goat anti-rabbit secondary antibody conjugated with Tritc (Santa Cruz Biotechnology) were used as needed. After the nuclei were counterstained with 4',6-diamidino-2-phenylindole (DAPI; Sigma), immunofluorescence images were visualised and recorded using an Olympus microscope system X51 or Olympus LSCM FV1000.

### Western blotting

One well of a 24-well plate of 60-day-old differentiated cells was lysed with RIPA lysis buffer (Biomiga) for western blotting. Blots were incubated with mouse anti-human CTNT, mouse anti-human  $\beta$ -MHC, rabbit anti-human MLC-2V, goat anti-ANF, mouse anti-human MLC-2A, mouse anti-human  $\beta$ -actin, rabbit anti-phospho smad1/5/8 and rabbit anti-smad1/5/8 separately. Blots were then incubated with HRP-conjugated goat anti-mouse or anti-rabbit antibody.

### Acknowledgments

We thank Drs Joy Fleming and Juanjuan Feng of the Institute of Biophysics, and Dr Paul Bornstein of the University of Washington for critical discussion and revision of the manuscript, and Dr Shigang He of the Institute of Biophysics for support and access to patch clamp technology. This work was supported by the Hi-Tech Research and Development Program of China (863 Program) (2006AA02A106), the National Basic Research Program of China (2006CB943901, 2010CB945024, and 2011CB965002), the Knowledge Innovation Program of the Chinese Academy of Sciences (KSCX2-YW-R-50), and the National Foundation of Science and Technology (30640005).

### References

- 1 He JQ, Ma Y, Lee Y, Thomson JA, Kamp TJ. Human embryonic stem cells develop into multiple types of cardiac myocytes: action potential characterization. *Circ Res* 2003; **93**:32-39.
- 2 Laflamme MA, Chen KY, Naumova AV, *et al.* Cardiomyo-

cytes derived from human embryonic stem cells in pro-survival factors enhance function of infarcted rat hearts. *Nat Biotechnol* 2007; **25**:1015-1024.

- 3 Yang L, Soonpaa MH, Adler ED, *et al.* Human cardiovascular progenitor cells develop from a KDR+ embryonic-stem-cell-derived population. *Nature* 2008; **453**:524-528.
- 4 Zhu WZ, Xie Y, Moyes KW, *et al.* Neuregulin/ErbB signaling regulates cardiac subtype specification in differentiating human embryonic stem cells. *Circ Res* 2010; **107**:776-786.
- 5 Kehat I, Khimovich L, Caspi O, *et al.* Electromechanical integration of cardiomyocytes derived from human embryonic stem cells. *Nat Biotechnol* 2004; **22**:1282-1289.
- 6 Chen HS, Kim C, Mercola M. Electrophysiological challenges of cell-based myocardial repair. *Circulation* 2009; **120**:2496-2508.
- 7 Tran TH, Wang X, Browne C, *et al.* Wnt3a-induced mesoderm formation and cardiomyogenesis in human embryonic stem cells. *Stem Cells* 2009; **27**: 1869-1878.
- 8 Reppel M, Igelmund P, Egert U, *et al.* Effect of cardioactive drugs on action potential generation and propagation in embryonic stem cell-derived cardiomyocytes. *Cell Physiol Biochem* 2007; **19**:213-224.
- 9 Yuasa S, Itabashi Y, Koshimizu U, *et al.* Transient inhibition of BMP signaling by Noggin induces cardiomyocyte differentiation of mouse embryonic stem cells. *Nat Biotechnol* 2005; **23**:607-611.
- 10 Marvin MJ, Di Rocco G, Gardiner A, Bush SM, Lassar AB. Inhibition of Wnt activity induces heart formation from posterior mesoderm. *Genes Dev* 2001; **15**:316-327.
- 11 Schneider VA, Mercola M. Wnt antagonism initiates cardiogenesis in *Xenopus laevis*. *Genes Dev* 2001; **15**:304-315.
- 12 Korol O, Gupta RW, Mercola M. A novel activity of the Dickkopf-1 amino terminal domain promotes axial and heart development independently of canonical Wnt inhibition. *Dev Biol* 2008; **324**:131-138.
- 13 Xu RH, Chen X, Li DS, *et al.* BMP4 initiates human embryonic stem cell differentiation to trophoblast. *Nat Biotechnol* 2002; **20**:1261-1264.
- 14 Zhang P, Li J, Tan Z, *et al.* Short-term BMP-4 treatment initiates mesoderm induction in human embryonic stem cells. *Blood* 2008; **111**:1933-1941.
- 15 Schulze GE, Clay RJ, Mezza LE, *et al.* BMS-189453, a novel retinoid receptor antagonist, is a potent testicular toxin. *Toxicol Sci* 2001; **59**:297-308.
- 16 Keegan BR, Feldman JL, Begemann G, Ingham PW, Yelon D. Retinoic acid signaling restricts the cardiac progenitor pool. *Science* 2005; **307**:247-249.
- 17 Xavier-Neto J, Rosenthal N, Silva FA, *et al.* Retinoid signaling and cardiac anteroposterior segmentation. *Genesis* 2001;

- 31:97-104.
- 18 Patwardhan V, Fernandez S, Montgomery M, Litvin J. The rostro-caudal position of cardiac myocytes affect their fate. *Dev Dyn* 2000; **218**:123-135.
  - 19 Orts-Llorca F J. Determination of heart polarity (arterio-venous axis) in the chicken embryo. *Roux Arch Entwicklungsmechanik* 1967; **113**:17.
  - 20 Yutzey K, Gannon M, Bader D. Diversification of cardiomyogenic cell lineages *in vitro*. *Dev Biol* 1995; **170**:531-541.
  - 21 Hochgreb T, Linhares VL, Menezes DC, *et al.* A caudorostral wave of RALDH2 conveys anteroposterior information to the cardiac field. *Development* 2003; **130**:5363-5374.
  - 22 Xavier-Neto J, Neville CM, Shapiro MD, *et al.* A retinoic acid-inducible transgenic marker of sino-atrial development in the mouse heart. *Development* 1999; **126**:2677-2687.
  - 23 Gassanov N, Er F, Zagidullin N, *et al.* Retinoid acid-induced effects on atrial and pacemaker cell differentiation and expression of cardiac ion channels. *Differentiation* 2008; **76**:971-980.
  - 24 Bao ZZ, Bruneau BG, Seidman JG, Seidman CE, Cepko CL. Regulation of chamber-specific gene expression in the developing heart by *Irx4*. *Science* 1999; **283**:1161-1164.
  - 25 Fu JD, Jiang P, Rushing S, *et al.*  $\text{Na}^+/\text{Ca}^{2+}$  exchanger is a determinant of excitation-contraction coupling in human embryonic stem cell-derived ventricular cardiomyocytes. *Stem Cells Dev* 2010; **19**:773-782.
  - 26 Maltsev VA, Rohwedel J, Hescheler J, Wobus AM. Embryonic stem cells differentiate *in vitro* into cardiomyocytes representing sinusnodal, atrial and ventricular cell types. *Mech Dev* 1993; **44**:41-50.
  - 27 Cheng H, Lederer WJ. Calcium sparks. *Physiol Rev* 2008; **88**:1491-1545.
  - 28 Woo SH, Cleemann L, Morad M. Spatiotemporal characteristics of junctional and nonjunctional focal  $\text{Ca}^{2+}$  release in rat atrial myocytes. *Circ Res* 2003; **92**:e1-e11.
  - 29 Cleemann L, Wang W, Morad M. Two-dimensional confocal images of organization, density, and gating of focal  $\text{Ca}^{2+}$  release sites in rat cardiac myocytes. *Proc Natl Acad Sci USA* 1998; **95**:10984-10989.
  - 30 Shimoji K, Yuasa S, Onizuka T, *et al.* G-CSF promotes the proliferation of developing cardiomyocytes *in vivo* and in derivation from ESCs and iPSCs. *Cell Stem Cell* 2010; **6**:227-237.
  - 31 Xu RH, Peck RM, Li DS, *et al.* Basic FGF and suppression of BMP signaling sustain undifferentiated proliferation of human ES cells. *Nat Methods* 2005; **2**:185-190.
  - 32 Ying QL, Nichols J, Chambers I, Smith A. BMP induction of *Id* proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell* 2003; **115**:281-292.
  - 33 Wobus AM, Kaomei G, Shan J, *et al.* Retinoic acid accelerates embryonic stem cell-derived cardiac differentiation and enhances development of ventricular cardiomyocytes. *J Mol Cell Cardiol* 1997; **29**:1525-1539.
  - 34 Domian IJ, Chiravuri M, van der Meer P, *et al.* Generation of functional ventricular heart muscle from mouse ventricular progenitor cells. *Science* 2009; **326**:426-429.
  - 35 Takahashi K, Tanabe K, Ohnuki M, *et al.* Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007; **131**:861-872.
  - 36 Yu J, Vodyanik MA, Smuga-Otto K, *et al.* Induced pluripotent stem cell lines derived from human somatic cells. *Science* 2007; **318**:1917-1920.
  - 37 Xu C, Inokuma MS, Denham J, *et al.* Feeder-free growth of undifferentiated human embryonic stem cells. *Nat Biotech* 2001; **19**:971-974.
  - 38 Picht E, Zima AV, Blatter LA, Bers DM. SparkMaster: automated calcium spark analysis with ImageJ. *Am J Physiol Cell Physiol* 2007; **293**:C1073-C1081.

(Supplementary information is linked to the online version of the paper on the *Cell Research* website.)